

Hi Sheldon:

Thank you for sharing.

Gilbert: “Possible. . . But New Horizons was launched 9 years ago. The elites keep track of Wormwood/PX via the ESO in Chile and the South Pole Telescope. With those two locations they can track Wormwood perfectly in real time. Both locations give good astronomical "viewing" in the Infrared part of the spectrum.”

Terral: “There is no viewing the Black Star in the infrared spectrum using a ground-based telescope, or my resident astronomer (Don) could also make observations using his observatory-class telescope and IR accessories. We know the Black Star location and also know we cannot see it.”

Gilbert: “With its helium bolometers (Infrared receivers), the South Pole Telescope can see the HEAT coming off of a methane ice cube (at - 296.4 Degrees F). So you KNOW they can see Wormwood/PX with a surface temp of 700+ Degrees F. There is NO WAY they are NOT going to see something that hot and that big (the size of Jupiter) closing in on the solar system from the southern hemisphere sky. It's just NOT going to happen. They know precisely where it is at all times. They just don't want the public to know that it's headed this way.”

Terral: “This is total BS. The Black Star is likely between 5 and 10 kilometers in diameter with a mass 25 to 50 percent of the Sun surrounded by an event horizon behaving more like a black hole than any regular celestial/cataloged object. This invisible/collapsed binary twin to our Sun sits inside an extremely deep/narrow gravity well where microlensing (gravitational lensing effect) is perfected bending visible light like a shroud becoming a cloak of invisibility. This notion that the Black Star is coming up from the southern hemisphere of our solar system is preposterous, as no seismic pattern would be present and we have evidence of a seismic pattern going back to 1965 (Mensur Omerbashich +6 Seismicity Paper to Cornell April 11, 2011). Scripture records the prophecies about the earthquakes, volcanoes, tsunamis, signs in the stars, etc., because there is a seismic pattern plainly visible representing the 'birth pangs' like those upon a woman with child and that comes before the 'destruction' that comes 'suddenly' (1Thes. 5:1-5), which happens with a springtime Black Star/Earth crossing event 'before' the Black Star reaches perihelion near the Sun. We know the Black Star perihelion location is near Venus Orbit Path, which is why Venus' rotation is backwards compared to the other planets from interaction with a gigantic outside force.

http://cdn.zmescience.com/wp-content/uploads/2013/01/binary_stars9.jpg

The Sun and Black Star are very near the same plane like most binary/trinary star systems, but our main sequence star is larger and the Black Star is the remnant of our Sun's binary twin with a wide elliptical orbit. The Black Star is between zero and -1 degrees below the ecliptic plane of our Sun making observation difficult from the northern hemisphere, which is the reason telescopes are positioned in the southern hemisphere and as near the South Pole as possible for looking across and under the debris on the plane. Earth is going through the first of two earth change lull periods right now until August 15, 2015 (started April 25, 2015 with the Nepal Quake at the alignment), because our

planet began moving away from the Black Star in the Virgo Constellation for the Black Star/Earth magnetic portal connection to lengthen causing the internal conduits to switch from active over to passive state. All of that will change after August 15th running to the beginning of December 2015, when the next earth change uptick period will see the earthquake/volcanic activity increase around the globe with focus on the US/Canada west coast. These kinds of patterns would not be present, if the Black Star were coming up from the southern hemisphere as Gilbert suggests. In fact, I can find no evidence anywhere to support his claims that appear nonsensical from the perspective of the science. For example:

One of the most-important events to take place in the last 5 years of my Project Black Star Investigation happened on March 12-13, 2012 with the Magnetopause Reversal Event.

https://www.dropbox.com/s/v15h4b2etcn9w11/iswa_download.gif?dl=0

This event took place one year and one day after Fukushima and a week before the Guerrero Quake Event from 3.20.2012, as 'both' of these events were predicted in advance by my research.

<http://www.nature.com/news/martian-mystery-cloud-defies-explanation-1.16924>

Tell me if this Martian Dust Storm defies explanation??? Well? Look at the date! That's right. March 12, 2012. Okay. Where was Mars on that date?

<https://www.dropbox.com/s/gnitnoix1jnfzor/Screenshot%202015-02-19%2011.20.03.png?dl=0>

Now. Maybe Sheldon or Gilbert or Andy Lloyd or James McCanney or Bill Broussard can explain how any celestial influence can possibly create environmental conditions in the inner solar system causing Earth magnetosphere to reverse for 28 hours 'and' cause a Martian Dust Storm on the 'dark side' of the planet, if that object is coming from below the Sun????? Anyone want to try and make that case? I think not! The influence that caused these strange events that defies explanation MUST come from the quadrant of space between the Sun and the Leo/Virgo Constellations! Period! That same influence caused the 8.8 Chile Quake from 2.27.2010, Fukushima, Guerrero and the 7.8 Nepal Quake that was also predicted by my Project Black Star Investigation on April 25, 2015 at the same time that the Kilauea Volcano magma levels in Hawaii crested from near Black Star proximity at the recent alignment. Now we are in the lull period that was also predicted and the coming uptick period will begin right on schedule around August 15, 2015.”

Gilbert:

“In addition to those instruments, they still have the Hubble, the Spitzer and will soon have the Webb space telescope. And if they have to, they can turn the WISE back on whenever they want to. It ran out of Hydrogen ice but it can still see perfectly in the Near Infrared by using the cold temps of space as a reference temperature. So with all those Infrared sensitive "eyes" available to watch Wormwood coming inbound, there is NO WAY they are going to miss it. And when it gets close to the ecliptic on the southern side, we will be able to see it as well. So fear not. . . we will have lots of warning that it is coming inbound long before it gets up close and personal with the Earth. GE”

Terral: “While the Black Star is indeed coming closer, nothing in the above commentary from Gilbert has anything to do with this case. ZERO. If Gilbert were anywhere near the right path, then his commentary would include these seismic events and the backside-alignment quake events going back to the 7.1 Christchurch Quake from 9.4.2010 and the 7.3 Fiji Quake from 9.15.2011 and the 7.3 Japan Quake from October 25, 2013 and the 7.3 Indonesian Quake from November 15, 2014, again, because there is a CLEAR seismic pattern (birth pangs) that the typical PlanetX/Nibiru/Wormwood research simply chooses to ignore.

Thanks again for sharing,”

Terral